

## Original Research Article

## Birth Weight, Postnatal Weight Gain, and Adult Body Composition in Five Low and Middle Income Countries

CHRISTOPHER W. KUZAWA,<sup>1,2\*</sup> PEDRO C. HALLAL,<sup>3</sup> LINDA ADAIR,<sup>4</sup> SANTOSH K. BHARGAVA,<sup>5</sup> CAROLINE H.D. FALL,<sup>6</sup> NANETTE LEE,<sup>7</sup> SHANE A. NORRIS,<sup>8</sup> CLIVE OSMOND,<sup>6</sup> MANUEL RAMIREZ-ZEA,<sup>9</sup> HARSHPAL SINGH SACHDEV,<sup>10</sup> ARYEH D. STEIN,<sup>11</sup> AND CESAR G. VICTORA<sup>3</sup>; COHORTS Group

<sup>1</sup>Department of Anthropology, Northwestern University, Evanston, Illinois

<sup>2</sup>Cells 2 Society: The Center for Social Disparities and Health at the Institute for Policy Research, Northwestern University, Evanston, Illinois

<sup>3</sup>Postgraduate Program in Epidemiology, Federal University of Pelotas, Brazil

<sup>4</sup>Department of Nutrition, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina

<sup>5</sup>Sunder Lal Jain Hospital, Delhi, India

<sup>6</sup>Lifecourse Epidemiology Unit, University of Southampton, Southampton, United Kingdom

<sup>7</sup>Office of Population Studies Foundation, University of San Carlos, Cebu, Philippines

<sup>8</sup>Department of Paediatrics, MRC/WITS Developmental Pathways for Health Research Unit, University of the Witwatersrand, Johannesburg, South Africa

<sup>9</sup>Institute of Nutrition of Central America and Panama—INCAP, Guatemala City, Guatemala

<sup>10</sup>Sitaram Bhartiya Institute of Science and Research, New Delhi, India

<sup>11</sup>Hubert Department of Global Health, Rollins School of Public Health, Emory University, Atlanta, Georgia

**Objectives:** To evaluate the associations between birth weight (BW), infancy, and childhood weight gain and adult body composition.

**Methods:** Subjects included participants of five birth cohort studies from low and middle income nations (Brazil, Guatemala, India, Philippines, and South Africa;  $n = 3432$ ). We modeled adult body composition as a function of BW and conditional weight gain (CW), representing changes in weight trajectory relative to peers, in three age intervals (0–12 months, 12–24 months, 24 months–mid childhood).

**Results:** In 34 of 36 site- and sex-specific models, regression coefficients associated with BW and CWs were higher for adult fat-free than for fat mass. The strength of coefficients predicting fat-free mass relative to those predicting fat mass was greatest for BW, intermediate for CWs through 24 months, and weaker thereafter. However, because fat masses were smaller and showed larger variances than fat-free masses, weaker relationships with fat mass still yielded modest but significant increases in adult % body fat (PBF). CW at 12 months and mid-childhood tended to be the strongest predictors of PBF, whereas BW was generally the weakest predictor of PBF. For most early growth measures, a 1 SD change predicted less than a 1% change in adult body fat, suggesting that any health impacts of early growth on changes in adult body composition are likely to be small in these cohorts.

**Conclusions:** BW and weight trajectories up to 24 months tend to be more strongly associated with adult fat-free mass than with fat mass, while weight trajectories in mid-childhood predict both fat mass and fat-free mass. *Am. J. Hum. Biol.* 24:5–13, 2012. © 2011 Wiley Periodicals, Inc.

Patterns of growth and weight gain during fetal life, infancy, and childhood predict adult metabolic and cardiovascular diseases as a result of changes in organ growth, metabolic function, and other biological pathways (Barker, 1990,2003; Barker, et al., 2009; Ben-Shlomo and Kuh, 2002). Postnatal changes in body composition represent an important candidate mechanism by which early experience might influence adult chronic disease risk (Eriksson et al., 2001; Kuh et al., 2002; Laitinen et al., 2004; McCarthy et al., 2007; Victora et al., 2007). Notably, birth weight (BW) is positively related to later lean mass, which could help explain why lower BW individuals are at increased risk for cardiovascular disease (Wells, 2007; Yajnik, 2002). Early growth-mediated changes in both the quantity and pattern of body fat deposition could also have effects on traits such as insulin resistance, blood pressure, and lipid metabolism and is thus understood as an important pathway linking early developmental processes with adult health (Wells, 2007).

In high-income countries, some studies report that rapid weight gain in the months immediately following birth predicts a more adverse body composition and metabolic risk profile in adulthood (Baird et al., 2005; Monteiro

COHORTS group members: Fernando C. Barros, Bernardo L. Horta, and Denise P. Gigante (Universidade Federal de Pelotas, Brazil), Daniel Lopez and Mathew Mainwaring (Department of Paediatrics, MRC Mineral Metabolism Research Unit, University of the Witwatersrand, Johannesburg, South Africa), Linda Richter (Human Sciences Research Council, Durban, South Africa), Judith Borja (Office of Population Studies Foundation, University of San Carlos, Cebu City, Philippines), Reynaldo Martorell (Hubert Department of Global Health, Rollins School of Public Health, Emory University, USA), Darren Dahly (University of Leeds, UK), Lakshmi Ramakrishnan, Nikhil Tandon (All-India Institute of Medical Sciences, New Delhi), Dorairaj Prabhakaran (Centre for the Control of Chronic Diseases, New Delhi), Siddharth Ramji (Maulana Azad Medical College, New Delhi), Anita Khalil (The Heart Centre, New Delhi), Poonima Prabhakaran, K Srinath Reddy (Public Health Foundation of India); SK Dey Biswas (Indian Council of Medical Research); Tarun Gera (Fortis Hospital, New Delhi).

Contract grant sponsors: Wellcome Trust (UK), US National Institutes of Health, US National Science Foundation, US National Center for Health Statistics, Indian Council of Medical Research, British Heart Foundation, Human Sciences Research Council, South African Medical Research Council, the Mellon Foundation, South-African Netherlands Programme on Alternative Development, Anglo American Chairman's Fund, Fogarty International Center.

\*Correspondence to: Christopher W. Kuzawa, Department of Anthropology, Northwestern University, Evanston, IL 60208, USA.  
E-mail: kuzawa@northwestern.edu

Received 3 July 2011; Accepted 14 September 2011

DOI 10.1002/ajhb.21227

Published online 28 November 2011 in Wiley Online Library (wileyonlinelibrary.com).

and Victora, 2005; Ong et al., 2000). In contrast, others find that rapid weight gain later in childhood, rather than in infancy, predicts a more rapid onset of body fat deposition, increased obesity risk, and elevated risk for cardiovascular diseases (Reilly et al., 2005).

It is presently unclear whether early growth trajectories have distinct relationships with adult body composition in lower or middle income countries, where intergenerational histories of undernutrition are coupled with often rapid nutritional, lifestyle, and weight change (Adair and Prentice, 2004; Yajnik, 2004). Past studies in several cohorts in these settings report evidence for stronger relationships between BW or infancy weight gain and adult lean mass than fat mass, with rapid weight gain in later childhood generally more strongly related to adult fat mass (Li et al., 2003; Sachdev et al., 2005; Victora et al., 2007). These findings support the promotion of infant and childhood growth in populations in which early life undernutrition, as manifest in growth faltering, is common (Victora et al., 2010).

The Consortium on Health Oriented Research in Transitional Societies (COHORTS) pools longitudinal data from five well-characterized long-term cohorts in middle or lower income populations to clarify the role of early life influences on a range of functional and health outcomes (Richter et al., in press). In this journal, we recently reported strongest relationships between infancy growth faltering and adult height deficits in these populations (Stein et al., 2010). Here, we expand on these analyses to clarify the importance of early weight trajectories as predictors of adult body composition in these five cohorts. We used a uniform analysis strategy across populations to investigate the importance of BW and weight trajectories during different intervals of infancy and childhood as predictors of adult fat and fat-free mass.

## MATERIALS AND METHODS

This study used data from five studies (Table 1) that are pooling resources and data as part of the COHORTS (Richter et al., in press; Victora et al., 2008): the 1982 Pelotas Birth Cohort (Brazil; Victora and Barros 2006), the Institute of Nutrition of Central America and Panama—INCAP Oriente Longitudinal Study (Guatemala; Stein et al., 2008); the New Delhi Birth Cohort Study (India; Bhargava et al., 2004); the Cebu Longitudinal Health and Nutrition Survey (Philippines; Adair et al., 2011); and the Birth to Twenty cohort (South Africa; Richter et al., 2004). The methods of each have been published in detail elsewhere (Victora et al., 2008). All studies were approved by an appropriate Ethics or Institutional Review Board.

### *Birth, infancy, and childhood anthropometry*

BW was measured by the research teams in Brazil, India, and Guatemala. In the Philippines, BW was measured by birth attendants provided with hanging scales for home births or was obtained from hospital records for hospital births. In South Africa, BW was obtained from birth records. In all studies, weights during infancy and childhood were measured by research teams. We identified four child ages at which weight was measured in all five cohorts, namely birth, 12 months, 24 months, and an age we denote as mid-childhood, which varies across cohorts

TABLE 1. Overview of the five studies included in the analysis and description of body composition measurements

Cohort country (name)	Design	Cohort inception (year)	Adult follow-up (year)	Number included in analysis (n)	Initial cohort	Body composition measurement
Guatemala (INCAP)	Community intervention	1969–1977	2004	205	Intervention trial of high-energy and protein supplement in women and children <7 years in 1969 and born during 1969–1977 in four villages.	Hydrostatic-weighting validated equation using weight, height, and abdominal circumference
India (Delhi)	Prospective cohort	1969–1972	1998–2002	936	Babies born to a population of married women living in a defined area of Delhi. Primarily middle-class sample.	Predictive equation using triceps and subscapular skinfolds
Philippines (CLHNS)	Prospective cohort	1983–1984	2005	1,612	Pregnant women living in 33 randomly selected neighborhoods; 75% urban, all social classes.	Predictive equation using triceps and subscapular skinfolds
South Africa (Birth-to-20)	Prospective cohort	1990	2005	302	Babies born to pregnant women living in defined urban area, primarily poor; black sample.	Dual X-ray Absorptiometry (DEXA).
Brazil (Pelotas)	Prospective cohort	1982	2005	377 (males only)	99% of all births in City's maternity hospital in 1982. All social classes.	Deuterium-validated equation using bioelectrical impedance. Only available for males.

(48 months in Brazil, Guatemala and India, 60 months in South Africa, and 102 months in Philippines). In India, weights corresponding to the age intervals of the study were interpolated using individual weight curves. We computed age and sex-specific *Z*-scores using the WHO Growth Standards (2006). We interpolated weight at 48 months in the samples from South Africa and Philippines by linear interpolation of the WAZ at 24 months and mid-childhood and back-transformation of the WAZ at 48 months to the absolute weight.

#### Conditional weight

Weight estimates measured at different ages are strongly correlated, leading to problems of collinearity if models simultaneously evaluate body size at different ages as predictors of adult outcomes. Therefore, we modeled relationships between early weight and adult outcomes using conditional weights (CWs; Adair et al., 2009). For each time point in infancy and childhood, CW was calculated as the residual from site- and sex-stratified linear regression of weight (kg) at that age on BW and any prior postnatal weights. The models were also adjusted for exact age at the time of measurement and a squared age term to account for any nonlinearities. As residuals, the CWs are uncorrelated with all prior weight measures. When entered into a multiple regression including BW and all prior CW estimates, CW may be interpreted as the deviation in the preceding growth interval from the weight predicted by BW, any prior CWs, and the population- and sex-specific mean growth.

#### Adult anthropometry and body composition

Adult height and weight were obtained by research teams using standard methods. Adult body fat and fat-free mass were calculated using site-specific methods. In Brazil, bioelectrical impedance was measured, and these results were corrected on the basis of a validation study that used isotopic methods (Wells et al., 2003); these data are only available for males. In Guatemala, weight, height, and abdominal circumference were obtained and entered into a hydrostatic-weighing validated equation (Ramirez-Zea et al., 2006). The India and Philippines cohorts used published equations (Durnin and Womersley, 1974) that have been validated for use in Asian populations for estimating body fat from skinfold measures (Deurenberg et al., 2002); South Africa used dual X-ray absorptiometry (Hologic Delphi). Fat mass was calculated as % body fat (PBF)  $\times$  body weight, and fat-free mass as weight – fat mass. Pregnant women were excluded from all analyses. The body mass index (BMI) was calculated as  $\text{kg/m}^2$ , which was used to define overweight (BMI  $\geq 25 \text{ kg/m}^2$ ) and obesity (BMI  $\geq 30 \text{ kg/m}^2$ ). As the South African sample was less than 18 years of age, in this cohort overweight and obesity status were defined using the *z*bmica command in Stata, which defines overweight and obesity status using age- and sex-specific cut points recommended by the Childhood Obesity Working Group of the International Obesity Taskforce (Cole et al., 2000).

#### Control variables

Age at adult measurement was entered as a continuous variable. In females, we adjusted for parity—including prior live births (Brazil, India, and Guatemala) or all prior

pregnancies, including stillbirths, live births, and premature terminations (Philippines). None of the participants in the South Africa cohort had given birth at the time of body composition measurement. Height was included in the models predicting fat and fat-free mass.

#### Statistical analysis

All statistical analyses were conducted using Stata (version 10, College Station, TX). Subsamples with all necessary variables were defined and descriptive statistics were calculated. In constructing models, we first tested to see if key relationships varied significantly by gender and across site. There was significant heterogeneity by gender for models predicting fat mass ( $F_{4,3421} = 20.15, P < 0.00001$ ) and fat-free mass ( $F_{4,3421} = 33.20, P < 0.00001$ ). In gender-stratified models, there was significant heterogeneity among males across sites in relationships between early growth measures and both fat ( $F_{16,2051} = 1.94, P < 0.014$ ) and fat-free mass ( $F_{16,2051} = 4.93, P < 0.00001$ ), and among females for fat ( $F_{12,1329} = 3.74, P < 0.00001$ ) and fat-free mass ( $F_{12,1329} = 4.80, P < 0.00001$ ). Therefore, all models reported here were run separately by gender and site. In analyses predicting fat mass, two models are presented: the first is unadjusted, and the second adjusted for age and height (and in females, also parity).

A central question addressed in this article is whether early weight gain during different intervals of infancy and childhood has different relationships with adult fat-free and fat mass. We used multivariate regression models to simultaneously predict fat-free mass and fat mass from BW and CWs at 12, 24 months, and mid-childhood. All predictors (BW and CW scores) and outcomes (fat-free mass and fat mass) were run as site- and gender-specific standard deviation scores. Thus, the coefficients from these models may be interpreted as partial correlations, allowing direct comparison of the relative strength of relationships between each predictor and outcome and between coefficients calculated at different sites. For each predictor, postestimation tests were used to evaluate whether the regression coefficients predicting adult fat-free and fat mass differed. Finally, we ran comparable site- and sex-stratified models predicting adult PBF as an outcome.

## RESULTS

Early life and adult characteristics of the study sample are summarized in Table 2. With the exception of Brazil, BWs were roughly between 0.5 and 1.0 SD below the WHO norm. Average weight-for-age *Z*-score declined between birth and 24 months in Guatemala and India and between birth and mid-childhood in the Philippines, with comparably stable WAZ scores between birth and mid-childhood in South Africa and Brazil. Mean adult height among the males ranged between 1.63 m in Guatemala and Philippines to 1.74 m in Brazil, and among females, between 1.51 m in Guatemala and Philippines to 1.59 m in South Africa. As adults, the Philippines sample was the leanest as indicated by the BMI, with less than 10% of males and females classified as overweight or obese. At the other extreme, 58% of the Guatemalan females in the sample were classified as overweight or obese.

With few exceptions, CW tended to be positively related to adult fat mass at all ages and sites and in both genders,

TABLE 2. Descriptive characteristics of cohorts during early life and at adult follow-up<sup>a</sup>

	Guatemala		India		Philippines		South Africa		Brazil Male (n = 377)
	Male (n = 114)	Female (n = 91)	Male (n = 540)	Female (n = 396)	Male (n = 887)	Female (n = 725)	Male (n = 160)	Female (n = 142)	
Weight (kg)									
Birth	3.1 (0.6)	3.0 (0.4)	2.9 (0.5)	2.8 (0.4)	3.0 (0.4)	3.0 (0.4)	3.2 (0.5)	3.0 (0.5)	3.3 (0.5)
12 months	7.9 (1.1)	7.7 (1.0)	8.5 (1.1)	7.9 (1.1)	8.3 (1.0)	7.7 (0.9)	9.7 (1.4)	9.1 (1.3)	9.8 (1.2)
24 months	9.9 (1.1)	9.6 (1.0)	10.4 (1.3)	9.7 (1.2)	10.1 (1.1)	9.5 (1.1)	11.7 (1.7)	11.2 (1.6)	12.3 (1.4)
Mid-childhood <sup>b</sup>	13.8 (1.4)	13.4 (1.4)	14.0 (1.6)	13.2 (1.6)	13.2 (1.8)	12.9 (1.8)	16.2 (1.8)	15.6 (1.8)	16.7 (2.4)
WAZ <sup>c</sup>									
Birth	-0.59 (1.22)	-0.53 (0.97)	-1.03 (1.00)	-1.07 (0.90)	-0.71 (0.93)	-0.57 (0.97)	-0.45 (1.17)	-0.69 (1.12)	-0.06 (1.15)
12 months	-1.83 (1.20)	-1.31 (1.06)	-1.20 (1.14)	-1.16 (1.14)	-1.47 (1.08)	-1.36 (1.03)	-0.09 (1.31)	0.06 (1.09)	0.002 (1.13)
24 months	-1.89 (1.01)	-1.53 (0.89)	-1.45 (1.05)	-1.44 (1.07)	-1.67 (0.97)	-1.68 (1.00)	-0.56 (1.30)	-0.41 (1.24)	-0.12 (1.07)
Mid-childhood <sup>b</sup>	-1.37 (0.80)	-1.41 (0.77)	-1.27 (0.89)	-1.52 (0.91)	-2.00 (1.04)	-1.85 (0.93)	-0.21 (0.96)	-0.31 (0.93)	0.06 (1.18)
Adult									
Age (years)	31.4 (1.2)	31.2 (1.2)	29.4 (1.3)	29.4 (1.3)	21.5 (0.3)	21.5 (0.3)	15.6 (0.3)	15.6 (0.2)	23.1 (0.3)
Height (cm)	162.8 (6.2)	151.4 (5.2)	169.5 (6.1)	155.1 (5.7)	163.1 (5.8)	151.1 (5.4)	166.5 (7.7)	158.8 (6.5)	173.8 (6.4)
Weight (kg)	63.7 (10.4)	60.6 (10.0)	72.0 (14.0)	59.5 (13.2)	56.0 (9.3)	46.4 (8.2)	55.6 (12.1)	54.6 (10.1)	74.0 (15.2)
Fat-free mass (kg)	51.0 (5.4)	39.1 (3.1)	53.9 (8.0)	38.4 (5.8)	46.4 (5.7)	31.0 (4.2)	45.9 (6.8)	36.5 (4.6)	61.1 (10.2)
Fat mass (kg)	12.7 (6.1)	21.4 (7.3)	18.1 (6.9)	21.1 (8.0)	9.7 (4.5)	15.4 (4.7)	9.8 (7.9)	18.2 (7.1)	12.9 (5.6)
% Body fat	19.2 (6.1)	34.4 (6.0)	24.4 (5.7)	34.3 (7.0)	16.7 (5.1)	32.7 (4.8)	16.4 (8.3)	32.3 (7.0)	16.8 (3.9)
BMI (kg/m <sup>2</sup> )	24.0 (3.4)	26.4 (4.0)	25.0 (4.3)	24.6 (5.0)	21.0 (3.0)	20.3 (3.2)	20.0 (3.7)	21.7 (3.8)	24.5 (4.6)
Overweight <sup>d</sup>	26%	40%	39%	34%	8%	7%	7%	14%	29%
Obese <sup>d</sup>	7%	18%	10%	13%	2%	1%	4%	6%	10%
Parous		85%		75%		31%		0%	

<sup>a</sup>Mean (SD) unless otherwise noted (see Methods for references to published profiles of each cohort).

<sup>b</sup>48 months in Brazil, Guatemala and India, 60 months in South Africa, 102 months in Philippines; South Africa and Philippine data interpolated to 48 months (see methods).

<sup>c</sup>Weight-for-age Z-score (WHO, 2006).

<sup>d</sup>25 kg/m<sup>2</sup> ≤ BMI < 30 kg/m<sup>2</sup>, BMI ≥ 30 kg/m<sup>2</sup>, For South Africa: International Obesity Task Force cut points (see Methods).

TABLE 3. Coefficients from multiple regression models relating standardized conditional weight scores to adult fat mass (kg)

	Males						Females					
	Unadjusted			Adjusted <sup>a</sup>			Unadjusted			Adjusted <sup>a</sup>		
	β	95% CI	P	β	95% CI	P	β	95% CI	P	β	95% CI	P
Guatemala												
Birth	0.70	(-0.42, 1.82)	0.22	0.41	(-0.90, 1.71)	0.54	1.86	(0.37, 3.35)	0.02	1.62	(0.06, 3.18)	0.05
12 months	1.04	(-0.13, 2.21)	0.09	0.90	(-0.34, 2.13)	0.15	1.16	(-0.33, 2.65)	0.13	0.53	(-1.20, 2.26)	0.55
24 months	0.53	(-0.57, 1.63)	0.34	0.37	(-0.81, 1.55)	0.54	1.29	(-0.18, 2.76)	0.09	0.87	(-0.75, 2.48)	0.29
Mid-childhood	0.91	(-0.26, 2.07)	0.13	0.82	(-0.38, 2.02)	0.18	0.74	(-0.84, 2.33)	0.36	0.52	(-1.12, 2.16)	0.54
R <sup>2</sup>		0.07			0.07			0.11			0.17	
India												
Birth	1.05	(0.52, 1.58)	0.0001	0.74	(0.23, 1.26)	0.005	1.76	(1.04, 2.48)	0.0001	1.58	(0.85, 2.32)	0.0001
12 months	1.79	(1.26, 2.33)	0.0001	1.56	(1.00, 2.12)	0.0001	2.37	(1.64, 3.10)	0.0001	2.00	(1.20, 2.80)	0.0001
24 months	1.29	(0.75, 1.83)	0.0001	1.22	(0.70, 1.74)	0.0001	1.10	(0.38, 1.81)	0.003	0.92	(0.20, 1.64)	0.02
Mid-childhood	1.51	(0.98, 2.05)	0.0001	1.52	(1.01, 2.04)	0.0001	1.23	(0.50, 1.97)	0.001	1.28	(0.54, 2.02)	0.001
R <sup>2</sup>		0.17			0.29			0.19			0.23	
Philippines												
Birth	0.35	(0.11, 0.60)	0.004	0.43	(0.18, 0.68)	0.001	0.81	(0.53, 1.08)	0.0001	0.82	(0.53, 1.10)	0.0001
12 months	1.73	(1.46, 1.99)	0.0001	1.89	(1.60, 2.18)	0.0001	1.37	(1.07, 1.66)	0.0001	1.40	(1.07, 1.72)	0.0001
24 months	0.59	(0.34, 0.84)	0.0001	0.65	(0.39, 0.91)	0.0001	0.46	(0.18, 0.75)	0.002	0.40	(0.11, 0.70)	0.008
Mid-childhood <sup>b</sup>	1.91	(1.67, 2.16)	0.0001	2.03	(1.77, 2.28)	0.0001	2.25	(1.98, 2.53)	0.0001	2.26	(1.97, 2.55)	0.0001
R <sup>2</sup>		0.33			0.34			0.35			0.36	
South Africa												
Birth	1.69	(0.54, 2.84)	0.004	1.89	(0.68, 3.10)	0.002	0.56	(-0.44, 1.56)	0.272	1.07	(0.05, 2.09)	0.04
12 months	1.61	(0.40, 2.82)	0.009	1.86	(0.56, 3.16)	0.005	3.10	(2.07, 4.13)	0.0001	3.62	(2.57, 4.67)	0.0001
24 months	0.76	(-0.38, 1.90)	0.19	0.87	(-0.30, 2.03)	0.145	0.43	(-0.56, 1.42)	0.40	0.78	(-0.18, 1.75)	0.12
Mid-childhood <sup>b</sup>	2.07	(0.94, 3.19)	0.0001	2.29	(1.09, 3.50)	0.0001	2.50	(1.46, 3.54)	0.0001	3.17	(2.10, 4.23)	0.0001
R <sup>2</sup>		0.16			0.17			0.30			0.36	
Brazil												
Birth	0.98	(0.51, 1.45)	0.0001	0.92	(0.43, 1.42)	0.0001						
12 months	1.51	(1.02, 2.00)	0.0001	1.45	(0.93, 1.97)	0.0001						
24 months	1.49	(1.00, 1.97)	0.0001	1.43	(0.93, 1.94)	0.0001						
Mid-childhood	2.16	(1.70, 2.62)	0.0001	2.16	(1.70, 2.62)	0.0001						
R <sup>2</sup>		0.33			0.34							

<sup>a</sup>Adjusted for age, height and in females also for parity.

<sup>b</sup>Interpolated (see Methods).

although there was heterogeneity in these relationships (Table 3). Among males, CW at 12 months and in mid-childhood were strongest predictors of adult fat mass in

Brazil, Guatemala, India, and the Philippines with CW at 48 months strongest in South Africa. In females, a similar pattern was present for the Philippines and South Africa

TABLE 4. Coefficients from multiple regression models relating standardized conditional weight scores to adult fat-free mass (kg)

	Males						Females					
	Unadjusted			Adjusted <sup>a</sup>			Unadjusted			Adjusted <sup>a</sup>		
	$\beta$	95% CI	<i>P</i>	$\beta$	95% CI	<i>p</i>	$\beta$	95% CI	<i>p</i>	$\beta$	95% CI	<i>p</i>
Guatemala												
Birth	2.31	(1.48, 3.15)	0.0001	0.90	(0.08, 1.72)	0.031	1.13	(0.55, 1.70)	0.0001	0.75	(0.22, 1.28)	0.006
12 months	1.34	(0.47, 2.22)	0.003	0.63	(-0.15, 1.40)	0.112	1.02	(0.44, 1.59)	0.001	0.33	(-0.26, .92)	0.274
24 months	1.48	(0.66, 2.30)	0.001	0.67	(-0.07, 1.41)	0.075	0.72	(0.15, 1.28)	0.013	0.26	(-0.29, 0.82)	0.343
Mid-childhood	1.19	(0.31, 2.06)	0.008	0.80	(0.04, 1.55)	0.038	0.50	(-0.11, 1.11)	0.107	0.26	(-0.30, 0.82)	0.363
<i>R</i> <sup>2</sup>		0.34			0.54		0.28				0.47	
India												
Birth	1.65	(1.08, 2.22)	0.0001	1.03	(0.47, 1.58)	0.0001	1.62	(1.15, 2.09)	0.0001	1.20	(0.73, 1.66)	0.0001
12 months	3.21	(2.64, 3.78)	0.0001	2.12	(1.52, 2.73)	0.0001	2.18	(1.70, 2.66)	0.0001	1.43	(0.92, 1.94)	0.0001
24 months	1.79	(1.22, 2.37)	0.0001	1.28	(0.72, 1.84)	0.0001	1.35	(0.88, 1.82)	0.0001	1.02	(0.56, 1.48)	0.0001
Mid-childhood	2.04	(1.47, 2.61)	0.0001	1.55	(0.99, 2.10)	0.0001	1.08	(0.59, 1.56)	0.0001	0.89	(0.42, 1.36)	0.0001
<i>R</i> <sup>2</sup>		0.31			0.39		0.32				0.40	
Philippines												
Birth	1.21	(0.95, 1.47)	0.0001	0.90	(0.65, 1.16)	0.0001	1.14	(0.92, 1.37)	0.0001	0.90	(0.68, 1.13)	0.0001
12 months	2.83	(2.55, 3.11)	0.0001	2.18	(1.88, 2.47)	0.0001	1.38	(1.14, 1.62)	0.0001	1.02	(0.77, 1.28)	0.0001
24 months	1.25	(0.98, 1.52)	0.0001	0.95	(0.68, 1.21)	0.0001	0.61	(0.37, 0.84)	0.0001	0.36	(0.13, 0.59)	0.002
Mid-childhood <sup>b</sup>	2.74	(2.48, 3.01)	0.0001	2.31	(2.04, 2.57)	0.0001	2.08	(1.86, 2.31)	0.0001	1.80	(1.57, 2.03)	0.0001
<i>R</i> <sup>2</sup>		0.52			0.57		0.44				0.51	
South Africa												
Birth	2.05	(1.20, 2.89)	0.0001	1.04	(0.35, 1.73)	0.004	1.80	(1.30, 2.30)	0.0001	1.38	(0.89, 1.88)	0.0001
12 months	2.32	(1.44, 3.21)	0.0001	1.12	(0.38, 1.87)	0.003	2.11	(1.59, 2.62)	0.0001	1.66	(1.16, 2.17)	0.0001
24 months	0.82	(-0.02, 1.65)	0.056	0.38	(-0.28, 1.05)	0.259	1.29	(0.80, 1.78)	0.0001	1.09	(0.63, 1.56)	0.0001
Mid-childhood <sup>b</sup>	2.94	(2.11, 3.77)	0.0001	1.84	(1.15, 2.53)	0.0001	1.75	(1.23, 2.27)	0.0001	1.32	(0.81, 1.84)	0.0001
<i>R</i> <sup>2</sup>		0.39			0.63		0.57				0.64	
Brazil												
Birth	2.70	(1.89, 3.52)	0.0001	2.04	(1.20, 2.87)	0.0001						
12 months	2.97	(2.12, 3.83)	0.0001	2.32	(1.45, 3.19)	0.0001						
24 months	3.17	(2.32, 4.01)	0.0001	2.57	(1.71, 3.42)	0.0001						
Mid-childhood	3.54	(2.75, 4.34)	0.0001	3.41	(2.63, 4.19)	0.0001						
<i>R</i> <sup>2</sup>		0.39			0.42							

<sup>a</sup>Adjusted for age, height and in females also for parity.

<sup>b</sup>Interpolated (see Methods).

but not Guatemala or India where BW or CW at 12 months tended to be strongest predictors of fat mass. These relationships were largely unchanged after adjustment for height and age (and in female's parity). CW scores tended to relate to fat-free mass in a similar pattern across sites and genders (Table 4). However, the strength of relationships varied by outcome and gender. Comparing findings from Tables 3 and 4, each SD-score change in CW predicted roughly double the kg change in adult fat-free mass compared with the predicted change in adult fat mass among males, whereas coefficients predicting adult fat-free and fat mass were more comparable among females. Unlike the models predicting fat mass, coefficients relating CW scores with FFM were substantially attenuated in every site and in both genders after adjustment for adult height.

We next used multivariate regression to evaluate the relative strength of coefficients relating CW scores at different ages to model adult fat-free mass and fat mass simultaneously as outcomes. Fat mass and fat-free mass were modeled as site- and gender-specific SD-scores to allow direct comparison of their respective coefficients, which can be interpreted as partial correlations. In these models, the coefficients for fat-free mass were greater than those for fat mass in 34 of 36 comparisons (Table 5). In formal postestimation tests comparing the strength of relationships between each CW score and fat-free and fat mass, there was consistent evidence across sites and genders for stronger effects of BW on fat-free mass than on fat mass (in eight of nine models) and for no differential

effects of CW at mid-childhood on fat-free mass or fat mass (in eight of nine models). There was more site- and gender-specific heterogeneity in effects of CW measured at the intervening ages, with CW at both 12 and 24 months a stronger predictor of fat-free mass than fat mass in only four of nine models.

The multivariate models showed that nearly all BW or early CW measures were more strongly correlated with changes in fat-free mass than with fat mass, when these are expressed as *Z* scores. Fat masses tend to equal 1/3 to 1/5th of fat-free masses in our cohorts (Table 2), but fat mass (coefficients of variation of 10–15%) was more variable than fat-free mass (coefficients of variation around 30–70%). This implies that a relatively weaker effect on fat mass *Z* scores could still increase the percentage of adult body weight accounted for by fat. To test this possibility, a final set of models (Table 6) evaluated relationships between BW and CW scores and adult PBF adjusting for age and in females also parity. Consistent with this expectation, nearly all early growth measures predicted a positive change in PBF. Among males, CW at 12 months and mid-childhood CW tended to be strongest predictors of PBF at each site, with mid-childhood CW the strongest coefficient in all cohorts. Among all male cohorts but South Africa, BW was the weakest predictor of PBF. Among females, the Philippines and South Africa followed a pattern similar to that of the males, with PBF most strongly related to CW at 12 months and in mid-childhood. However, there was more heterogeneity across female cohorts: BW was the strongest predictor of PBF in Guate-

TABLE 5. Coefficients from multivariate models relating standardized conditional weight scores to adult fat and fat-free mass (SD units)<sup>a</sup>

	Males					Females				
	Fat mass		Fat-free mass		<i>p</i> -value <sup>b</sup>	Fat mass		Fat-free mass		<i>p</i> -value <sup>b</sup>
	$\beta$	95%CI	$\beta$	95%CI		$\beta$	95%CI	$\beta$	95%CI	
Guatemala										
Birth	0.12	(-0.07, 0.30)	0.43	(0.27, 0.59)	0.0001	0.27	(0.06, 0.47)	0.37	(0.18, 0.56)	0.1
12 months	0.17	(-0.02, 0.37)	0.26	(0.09, 0.42)	0.30	0.15	(-0.06, 0.36)	0.33	(0.14, 0.52)	0.007
24 months	0.09	(-0.09, 0.27)	0.28	(0.13, 0.43)	0.01	0.17	(-0.04, 0.38)	0.24	(0.05, 0.43)	0.28
Mid-childhood	0.15	(-0.05, 0.34)	0.21	(0.05, 0.38)	0.40	0.10	(-0.13, 0.32)	0.16	(-0.05, 0.36)	0.37
India										
Birth	0.13	(0.06, 0.21)	0.20	(0.13, 0.27)	0.02	0.22	(0.13, 0.31)	0.28	(0.20, 0.36)	0.04
12 months	0.27	(0.20, 0.35)	0.40	(0.33, 0.47)	0.0001	0.28	(0.19, 0.37)	0.36	(0.28, 0.45)	0.006
24 months	0.20	(0.13, 0.27)	0.23	(0.15, 0.30)	0.35	0.13	(0.04, 0.22)	0.23	(0.15, 0.31)	0.0005
Mid-childhood	0.24	(0.17, 0.32)	0.26	(0.19, 0.33)	0.59	0.17	(0.08, 0.26)	0.20	(0.11, 0.28)	0.40
Philippines										
Birth	0.08	(0.03, 0.13)	0.21	(0.16, 0.26)	0.0001	0.17	(0.12, 0.23)	0.28	(0.22, 0.33)	0.0002
12 months	0.39	(0.33, 0.44)	0.49	(0.44, 0.54)	0.0005	0.30	(0.24, 0.36)	0.34	(0.28, 0.40)	0.16
24 months	0.13	(0.07, 0.18)	0.22	(0.17, 0.27)	0.002	0.09	(0.02, 0.15)	0.13	(0.08, 0.19)	0.11
Mid-childhood <sup>c</sup>	0.43	(0.37, 0.48)	0.48	(0.43, 0.52)	0.09	0.49	(0.43, 0.55)	0.49	(0.44, 0.54)	0.91
South Africa										
Birth	0.21	(0.07, 0.36)	0.31	(0.18, 0.43)	0.31	0.07	(-0.07, 0.21)	0.39	(0.28, 0.50)	0.0001
12 months	0.21	(0.05, 0.36)	0.35	(0.22, 0.49)	0.12	0.43	(0.28, 0.57)	0.46	(0.34, 0.57)	0.74
24 months	0.10	(-0.05, 0.24)	0.13	(0.01, 0.26)	0.70	0.07	(-0.07, 0.21)	0.29	(0.18, 0.40)	0.006
Mid-childhood <sup>c</sup>	0.26	(0.12, 0.41)	0.44	(0.32, 0.56)	0.05	0.36	(0.21, 0.51)	0.39	(0.27, 0.50)	0.74
Brazil										
Birth	0.17	(0.09, 0.26)	0.26	(0.18, 0.34)	0.003					
12 months	0.27	(0.18, 0.35)	0.29	(0.20, 0.37)	0.49					
24 months	0.26	(0.18, 0.35)	0.31	(0.22, 0.39)	0.15					
Mid-childhood	0.39	(0.30, 0.47)	0.35	(0.27to 0.43)	0.25					

<sup>a</sup>Adjusted for age and in females also for parity.<sup>b</sup>Significance of difference in regression coefficients predicting fat-free and fat mass evaluated using multivariate regression.<sup>c</sup>Interpolated (see Methods).

mala, whereas in India PBF was most strongly related to BW and 12 month CW.

## DISCUSSION

We address an important public health problem using a uniform modeling approach and a common analytic strategy in five well-characterized cohorts representing different political-economic, cultural, and nutritional circumstances. Despite differences in economic and other conditions across populations and in methods used to measure body composition, we found certain consistencies across the five sites. We also found that results were contingent on the body composition measure employed. When adult body composition was modeled as separate compartments of fat-free and fat mass, BW tended to be a stronger predictor of adult fat-free mass than of fat mass irrespective of gender and site, consistent with prior evidence of beneficial effects of fetal growth on adult body composition (Singhal et al., 2003; Wells et al., 2007). In contrast, CW measured in mid-childhood was not differentially related to fat-free or fat mass in adulthood in any site or gender, pointing to fewer potential benefits to lean tissue in relation to weight gain at older ages. Modeling PBF as an outcome revealed that the correlations of most BW and CW scores with fat mass, although weaker than correlations with those with fat-free mass when both are expressed in standardized units, were sufficient to modestly increase PBF. In males, BW was generally the weakest predictor of PBF. Although there was heterogeneity across models, CW at 12 months, reflecting weight gain in the first year of life, and mid-childhood CW, reflecting early childhood weight gain, tended to be the strongest predictors of adult

TABLE 6. Coefficients from regression models relating standardized conditional weight scores to adult percent body fat<sup>a</sup>

	Males			Females		
	$\beta$	95% CI	<i>P</i>	$\beta$	95% CI	<i>p</i>
Guatemala						
Birth	0.01	(-1.13, 1.16)	0.981	1.46	(0.21, 2.70)	0.0220
12 months	0.90	(-0.30, 2.10)	0.14	0.59	(-0.69, 1.88)	0.3590
24 months	0.17	(-0.96, 1.29)	0.772	0.76	(-0.50, 2.02)	0.2350
Mid-childhood	1.00	(-0.20, 2.21)	0.102	0.51	(-0.84, 1.86)	0.4580
<i>R</i> <sup>2</sup>		0.05			0.14	
India						
Birth	0.45	(0.03, 0.88)	0.0370	1.04	(0.39, 1.68)	0.0020
12 months	0.88	(0.45, 1.31)	0.0001	1.58	(0.91, 2.24)	0.0001
24 months	0.82	(0.39, 1.26)	0.0001	0.49	(-0.16, 1.13)	0.1370
Mid-childhood	0.99	(0.56, 1.42)	0.0001	0.90	(0.23, 1.56)	0.0080
<i>R</i> <sup>2</sup>		0.22			0.15	
Philippines						
Birth	0.16	(-0.14, 0.47)	0.2940	0.31	(-0.01, 0.64)	0.0560
12 months	1.46	(1.13, 1.79)	0.0001	0.99	(0.65, 1.34)	0.0001
24 months	0.36	(0.04, 0.68)	0.0260	0.20	(-0.14, 0.54)	0.2600
Mid-childhood <sup>b</sup>	1.65	(1.34, 1.95)	0.0001	1.57	(1.25, 1.90)	0.0001
<i>R</i> <sup>2</sup>		0.18			0.15	
South Africa						
Birth	1.49	(0.22, 2.75)	0.021	-0.55	(-1.63, 0.53)	0.3160
12 months	1.17	(-0.16, 2.51)	0.085	2.19	(1.08, 3.30)	0.0001
24 months	0.67	(-0.59, 1.93)	0.296	-0.29	(-1.35, 0.78)	0.5950
Mid-childhood <sup>b</sup>	1.64	(0.39, 2.88)	0.01	1.92	(0.80, 3.04)	0.0001
<i>R</i> <sup>2</sup>		0.10			0.18	
Brazil						
Birth	0.40	(0.04, 0.75)	0.0290			
12 months	0.96	(0.59, 1.33)	0.0001			
24 months	0.81	(0.44, 1.18)	0.0001			
Mid-childhood	1.25	(0.90, 1.60)	0.0001			
<i>R</i> <sup>2</sup>		0.22				

<sup>a</sup>Adjusted for age and in females also for parity.<sup>b</sup>Interpolated (see Methods).

PBF in both genders. These findings show that patterns of early life weight gain predict adult body composition, while also documenting heterogeneity in these relationships by gender and across populations.

Modeling relationships between early weight gain and adult fatness requires grappling with several issues, including the choice of outcomes, the age ranges of early life weight gain to be investigated as predictors, and the related statistical problem that weight gain in a given age interval tends to be highly correlated with gain in adjacent intervals. With respect to the first issue, adult BMI is a commonly reported outcome in studies of early life determinants of adult body composition, with consistent positive associations being reported with BW and faster weight gains in infancy and childhood (Baird et al., 2005; Monteiro and Victora, 2005). Because BMI does not separate fat from fat-free mass, such results must be interpreted with caution. As an alternative approach, we investigated fat mass as an outcome, and evaluated whether early weight trajectories are associated more strongly with fat than with fat-free mass in adulthood. In our view, this is a useful approach, because one would like children to build up fat-free mass without gaining excess body fat as adults. Finally, we also considered the % of body weight that is fat as an outcome, which provides insight into how differential effects on lean and fat compartments contribute to changes in body composition.

The second set of issues involves how best to model early weight trajectories as predictors of adult body composition. Past work has shown that it is important to separate the long-term effects of early and later childhood weight gain (Victora et al., 2008). Some of the first studies on the long-term consequences of growth patterns in childhood reported on weight gains from birth to school age (Eriksson et al., 2001; Parsons et al., 2001) and were therefore unable to identify age ranges associated with different risks. We opted to model the relationship between early weight during different intervals from birth to mid-childhood, which allowed us to tease apart the relative importance of change in weight at different ages to adult fat and fat-free mass. However, because weight at the end of an age range is the same as the weight at the beginning of the next range, collinearity would be unavoidable if simple weight gains or weight velocities were modeled as independent predictors. To avoid this problem, we used CW scores, which estimate the effect of that component of weight gain in each interval that is independent of gain in prior intervals.

We found that associations relating early CW scores to adult body composition were not consistent for males and females, and in addition varied substantially across sites. We anticipated heterogeneity across sites in light of the many factors that differentiated the sites and that we were not able to fully control in our models. Notably, the age at which subjects had their adult body composition measured varied by site, as did the method used to estimate fat and fat-free mass. In addition, the populations varied in prevalence of low BW, early life undernutrition and adult overweight, reflecting diverse settings in which the nutrition and epidemiologic transitions have had differential influences on adult health and nutritional status. The significant heterogeneity in the pattern and magnitude of coefficients across site and by gender required that we pursue stratified rather than pooled models. Nevertheless, underlying consistencies were apparent across the

sites that point to the differential importance of weight gain in different early life age intervals as predictors of distinct components of adult body composition.

CW scores are positive when growth in the preceding interval was faster than expected based on that individual's weight trajectory entering that interval and the mean population growth during the interval. Thus, we interpret our CW models as indicating that faster than expected weight gains from birth to 12 months and from 24 months to mid-childhood tend to be more strongly associated with adult fat mass than are BW or relatively rapid weight gain from 12–24 months. Adjusting for height attenuated coefficients predicting fat-free mass, especially in males but left coefficients predicting fat mass largely unchanged. This suggests that early weight gain influences fat-free mass in part via effects on adult stature, consistent with prior analysis in these cohorts (Stein et al., 2010).

Certain consistencies emerged across sites when we modeled fat and fat-free masses simultaneously as outcomes. Here fat and fat-free mass were modeled as site- and sex-specific SD scores, thus allowing for a direct comparison of effect sizes of each CW score predicting fat and fat-free mass while also taking into account the average differences in adult fat-free mass between males and females. In these models, BW and to a lesser extent CW in the first 2 years of life tended to be more strongly associated with fat-free than with fat mass, but this differential effect on fat-free mass was absent in eight of nine models by mid-childhood.

The average adult fat-free mass tended to be three to five times the mass of fat in the cohorts considered here, while variability for fat mass was also larger. Thus, even relatively weaker relationships between early growth and adult fat mass than for fat-free mass could increase the proportion of adult weight accounted for by fat. Our models predicting PBF confirmed this expectation. With few exceptions, faster earlier growth at any interval predicted an increase in adult PBF in both males and females. Although the strength of these effects did vary by age, gender, and site, there were certain consistencies across many of the cohorts, and in general, findings mirrored trends seen in models predicting fat mass and fat-free mass. Just as CW at 12 months and mid-childhood tended to be strongest predictors of fat mass (Table 3), the weight gain represented in these intervals generally predicted the largest % increase in adult body fat. In the males, mid-childhood CW was the strongest predictor of PBF in all the cohorts, while BW was the weakest predictor of PBF among all male cohorts but South Africa. Among females, CW at 12 months and in mid-childhood tended to be strongest predictors of PBF.

Although many early growth measures were significant predictors of adult PBF, these effects were generally small. A full standard deviation change in most early growth measures, which represents a large change in early weight trajectory, predicted on average a 0.85% change in adult body fat. The health impacts of a change of this magnitude are likely to be negligible in most of our cohorts, many of whom are relatively lean. Based on these modest effect sizes, it seems likely that the many benefits to health and function known to result from improvements in early nutrition and growth (e.g., Victora et al., 2008) will outweigh any negative effects on health operating through changes in body composition.

Previously published single-site analyses of our cohorts are generally in line with the findings reported here, despite the fact that these analyses did not rely on CWs and none modeled fat and fat-free mass simultaneously. In Brazil, adult height was primarily determined by fetal and infant growth, whereas weight-related indices were more strongly influenced by later growth (Victora et al., 2007). In India, BMI changes in infancy and early childhood were correlated more strongly with adult lean mass than with adiposity or central adiposity, whereas BMI changes in late childhood or adolescence were associated with increased adult adiposity and central adiposity (Sachdev et al., 2005). In Guatemala, BMI in infancy and later childhood was positively associated with adult BMI, % fat, abdominal circumference, and fat-free mass; after the age of 3 years, associations were stronger with BMI, % fat, and abdominal circumference than with fat-free mass (Corvalan et al., 2007). In a recent analysis of the Philippine cohort, weight gain from birth to 6 months of age was found to be an especially strong predictor of adult stature, fat-free mass, and strength in males but these relationships were weaker in females (Kuzawa et al., 2010).

Other studies in high income populations have addressed these issues. Positive associations between BW and lean mass—to a greater extent than with fat mass—have been reported in the United Kingdom (Singhal et al., 2003) and from a younger cohort in Brazil (Wells et al., 2005). In addition, the finding that weight gain from 24 months to mid-childhood is a stronger predictor of adult adiposity than is earlier weight gains has been reported in cohorts from the United Kingdom (Ong et al., 2000), Brazil (Wells et al., 2005), and Sweden (Ekelund et al., 2007). At the same time, the relatively low average BW of most of our study populations underscores the distinct starting point of growth trajectories in these settings compared with most higher income populations. Future studies in high income nations using the analysis approach used here would help clarify how the predictors of adult body composition might vary across these contexts.

The different strengths of the coefficients linking CW during the first and second years of life with adult body composition may partly reflect absolute differences in weight gain during each growth interval or the increasing stability of growth trajectories as children mature. In our cohorts, absolute gains in body weight during the first year of life were roughly three times the gains in the second year of life (Table 2), suggesting a greater potential for variation in early postnatal growth to influence adult body composition. In addition, there may be important biological critical periods during specific developmental windows, such as the early postnatal surge in sex steroid production (Winter et al., 1976), which could influence body composition development and the pattern of gender differences in adult body composition (Kuzawa et al., 2010).

Our observational data can reveal patterns of association but cannot speak to causes. Indeed, it is possible that part of the association between early CW and adult body composition reflects the pleiotropic effects of genes. Polymorphic loci with effects on early weight growth patterns have been identified (Sovio et al., 2009), and if such genes also influence adult body composition, this could lead to correlations between early CW scores and adult body composition (Suarez et al., 1997). Although we cannot rule out such effects, prior research conducted in the cohorts analyzed here (Stein et al., 2010; Victora et al., 1987), and in

populations living under similar socioeconomic and political-economic settings (Billewicz and McGregor, 1982), has underscored the overwhelming importance of environmental factors such as nutrition, infant feeding practices, and early life infectious morbidity as determinants of skeletal growth and weight gain during the first 2–3 years of postnatal life, highlighting the potential for environmental contributions to the relationships that we document.

Additional limitations of the analyses reported here warrant mention. As alluded to above, methods for measuring adult body composition varied by site and some of the between-site heterogeneity undoubtedly reflects biases inherent in these methods. Although this limits the conclusions that we may draw regarding population differences, these concerns apply less to our findings of sex differences, which were found within site using common methods. The instruments used to measure diet, lifestyle, or environmental conditions across our five sites were also not directly comparable, which constrained our ability to adjust for potentially important covariates.

Although our analyses were limited by the ages at which weight measurements were available across sites, these measures allowed us to probe impacts at age intervals previously shown to be important as predictors of adult body composition in other populations. The ages of adult body composition measurement were more variable across sites, which likely contributed to the heterogeneity in relationships that we document. In this respect, the generally weaker relationships between early growth and fat-free mass among the South African males might partially reflect their relatively young age (~15.5 years), and the fact that males would have been at various stages of puberty and adult height attainment when body composition was measured. Future analysis of this cohort after they have achieved final adult stature would help clarify the importance of this source of biological variation as an influence on the relationships documented here. Although our focus here has been on body composition per se, future work should also evaluate associations with regional fat patterning, as this may be an important link between early growth processes and the metabolic syndrome and cardiovascular disease risk (Despres and Lemieux, 2006).

#### ACKNOWLEDGMENTS

Guatemala: Rafael Flores, Usha Ramakrishnan, Kathryn Yount, Ruben Grajeda, Paul Melgar, Humberto Mendez, Luis Fernando Ramirez, Jere Behrman, John Hoddinott, Agnes Quisumbing, Alexis Murphy, John Maluccio, Ann DiGirolamo; Philippines: Socorro Gultiano, Alan Feranil, Judith Borja, Josephine Avila, Lorna Perez, Barry Popkin; Brazil: Fernando Barros, Rosangela Lima, Jonathan Wells; India: Vinod Kapani, Rajeshwari Verma, Bhaskar Singh, Dileep Gupta; South Africa: John Pettifor, Noel Cameron, and Stella Fleetwood.

#### LITERATURE CITED

- Adair LS, Martorell R, Stein AD, Hallal PC, Sachdev HS, Prabhakaran D, Wills AK, Norris SA, Dahly DL, Lee NR, Victora CG. 2009. Size at birth, weight gain in infancy and childhood, and adult blood pressure in 5 low- and middle-income-country cohorts: when does weight gain matter? *Am J Clin Nutr* 89:1383–1392.
- Adair LS, Popkin BM, Akin JS, Guilkey DK, Gultiano S, Borja J, Perez L, Kuzawa CW, McDade T, Hindin MJ. 2011. Cohort Profile: The Cebu Longitudinal Health and Nutrition Survey. *Int J Epidemiol*. 40(3):619–25.

- Adair LS, Prentice AM. 2004. A critical evaluation of the fetal origins hypothesis and its implications for developing countries. *J Nutr* 134:191–193.
- Baird J, Fisher D, Lucas P, Kleijnen J, Roberts H, Law C. 2005. Being big or growing fast: systematic review of size and growth in infancy and later obesity. *BMJ* 331:929–934.
- Barker DJ. 1990. The fetal and infant origins of adult disease. *BMJ* 301:1111.
- Barker DJ. 2003. The developmental origins of adult disease. *Eur J Epidemiol* 18:733–736.
- Barker DJ, Osmond C, Kajantie E, Eriksson JG. 2009. Growth and chronic disease: findings in the Helsinki Birth Cohort. *Ann Hum Biol* 36:445–458.
- Ben-Shlomo Y, Kuh D. 2002. A life course approach to chronic disease epidemiology: conceptual models, empirical challenges and interdisciplinary perspectives. *Int J Epidemiol* 31:285–293.
- Bhargava SK, Sachdev HS, Fall CH, Osmond C, Lakshmy R, Barker DJ, Biswas SK, Ramji S, Prabhakaran D, Reddy KS. 2004. Relation of serial changes in childhood body-mass index to impaired glucose tolerance in young adulthood. *N Engl J Med* 350:865–875.
- Billewicz WZ, McGregor IA. 1982. A birth-to-maturity longitudinal study of heights and weights in two West African (Gambian) villages, 1951–1975. *Ann Hum Biol* 9:309–320.
- Cole TJM, Bellizzi C, Flegal KM, Dietz WH. 2000. Establishing a standard definition for child overweight and obesity worldwide: international survey. *Brit Med J* 320:1240–1243.
- Corvalan C, Gregory CO, Ramirez-Zea M, Martorell R, Stein AD. 2007. Size at birth, infant, early and later childhood growth and adult body composition: a prospective study in a stunted population. *Int J Epidemiol* 36:550–557.
- Despres JP, Lemieux I. 2006. Abdominal obesity and metabolic syndrome. *Nature* 444:881–887.
- Deurenberg P, Deurenberg-Yap M, Schouten FJ. 2002. Validity of total and segmental impedance measurements for prediction of body composition across ethnic population groups. *Eur J Clin Nutr* 56:214–220.
- Durnin JV, Womersley J. 1974. Body fat assessed from total body density and its estimation from skinfold thickness: measurements on 481 men and women aged from 16 to 72 years. *Br J Nutr* 32:77–97.
- Ekelund U, Ong KK, Linne Y, Neovius M, Brage S, Dunger DB, Wareham NJ, Rosner S. 2007. Association of weight gain in infancy and early childhood with metabolic risk in young adults. *J Clin Endocrinol Metab* 92:98–103.
- Eriksson J, Forsen T, Tuomilehto J, Osmond C, Barker D. 2001. Size at birth, childhood growth and obesity in adult life. *Int J Obes Relat Metab Disord* 25:735–740.
- Kuh D, Hardy R, Chaturvedi N, Wadsworth ME. 2002. Birth weight, childhood growth and abdominal obesity in adult life. *Int J Obes Relat Metab Disord* 26:40–47.
- Kuzawa CW, McDade TW, Adair LS, Lee N. 2010. Rapid weight gain after birth predicts life history and reproductive strategy in Filipino males. *Proc Natl Acad Sci USA* 107:16800–16805.
- Laitinen J, Pietilainen K, Wadsworth M, Sovio U, Jarvelin MR. 2004. Predictors of abdominal obesity among 31-year-old men and women born in Northern Finland in 1966. *Eur J Clin Nutr* 58:180–190.
- Li H, Stein AD, Barnhart HX, Ramakrishnan U, Martorell R. 2003. Associations between prenatal and postnatal growth and adult body size and composition. *Am J Clin Nutr* 77:1498–1505.
- McCarthy A, Hughes R, Tilling K, Davies D, Davey Smith G, Ben-Shlomo Y. 2007. Birth weight; postnatal, infant, and childhood growth; and obesity in young adulthood: evidence from the Barry Caerphilly Growth Study. *Am J Clin Nutr* 86:907–913.
- Monteiro PO, Victora CG. 2005. Rapid growth in infancy and childhood and obesity in later life—a systematic review. *Obes Rev* 6:143–154.
- Ong KK, Ahmed ML, Emmett PM, Preece MA, Dunger DB. 2000. Association between postnatal catch-up growth and obesity in childhood: prospective cohort study. *Brit Med J* 320:967–971.
- Parsons TJ, Power C, Manor O. 2001. Fetal and early life growth and body mass index from birth to early adulthood in 1958 British cohort: longitudinal study. *Brit Med J* 323:1331–1335.
- Ramirez-Zea M, Torun B, Martorell R, Stein AD. 2006. Anthropometric predictors of body fat as measured by hydrostatic weighing in Guatemalan adults. *Am J Clin Nutr* 83:795–802.
- Reilly JJ, Armstrong J, Dorosty AR, Emmett PM, Ness A, Rogers I, Steer C, Sherriff A. 2005. Early life risk factors for obesity in childhood: cohort study. *Brit Med J* 330:1357–1363.
- Richter LM, Norris SA, De Wet T. 2004. Transition from birth to ten to birth to twenty: the South African cohort reaches 13 years of age. *Paediatr Perinat Epidemiol* 18:290–301.
- Richter LM, Victora CG, Hallal PC, Adair LS, Bhargava SK, Fall CH, Lee N, Martorell R, Norris SA, Sachdev HS, Stein AD; COHORTS group. Cohort profile: the consortium of health-orientated research in transitioning societies. *Int J Epidemiol* (in press).
- Sachdev HS, Fall CH, Osmond C, Lakshmy R, Dey Biswas SK, Leary SD, Reddy KS, Barker DJ, Bhargava SK. 2005. Anthropometric indicators of body composition in young adults: relation to size at birth and serial measurements of body mass index in childhood in the New Delhi birth cohort. *Am J Clin Nutr* 82:456–466.
- Singhal A, Wells J, Cole TJ, Fewtrell M, Lucas A. 2003. Programming of lean body mass: a link between birth weight, obesity, and cardiovascular disease? *Am J Clin Nutr* 77:726–730.
- Sovio U, Bennett AJ, Millwood IY, Molitor J, O'Reilly PF, Timpson NJ, Kaakinen M, Laitinen J, Haukka J, Pillas D, Tzoulaki I, Molitor J, Hoggart C, Coin LJ, Whittaker J, Pouta A, Hartikainen AL, Freimer NB, Widen E, Peltonen L, Elliott P, McCarthy MI, Jarvelin MR. 2009. Genetic determinants of height growth assessed longitudinally from infancy to adulthood in the northern Finland birth cohort 1966. *PLoS Genet* 5:e1000409. doi: 10.1371/journal.pgen.1000409.
- Stein AD, Melgar P, Hoddinott J, Martorell R. 2008. Cohort profile: the Institute of Nutrition of Central America and Panama (INCAP) Nutrition Trial Cohort Study. *Int J Epidemiol* 37:716–720.
- Stein AD, Wang M, Martorell R, Norris SA, Adair LS, Bas I, Sachdev HS, Bhargava SK, Fall CH, Gigante DP, Victora CG; COHORTS group. 2010. Growth patterns in early childhood and final attained stature: data from five birth cohorts from low- and middle-income countries. *Am J Hum Biol* 22:353–359.
- Suarez F, Zeghoud F, Rossignol C, Walrant O, Garabedian M. 1997. Association between vitamin D receptor gene polymorphism and sex-dependent growth during the first two years of life. *J Clin Endocrinol Metab* 82:2966–2970.
- Victora CG, Adair L, Fall C, Hallal PC, Martorell R, Richter L, Sachdev HS. 2008. Maternal and child undernutrition: consequences for adult health and human capital. *Lancet* 371:340–357.
- Victora CG, Barros FC. 2006. Cohort profile: the 1982 Pelotas (Brazil) birth cohort study. *Int J Epidemiol* 35:237–242.
- Victora CG, Barros FC, Vaughan JP, Martines JC, Béria JU. 1987. Birth-weight, socio-economic status and growth of Brazilian infants. *Ann Hum Biol* 14:49–57.
- Victora CG, de Onis M, Hallal PC, Blossner M, Shrimpton R. 2010. Worldwide timing of growth faltering: revisiting implications for interventions. *Pediatrics* 125:e473–e480.
- Victora CG, Sibbritt D, Horta BL, Lima RC, Cole T, Wells J. 2007. Weight gain in childhood and body composition at 18 years of age in Brazilian males. *Acta Paediatr* 96:296–300.
- Wells JC. 2007. The programming effects of early growth. *Early Hum Dev* 83:743–748.
- Wells JC, Chomtho S, Fewtrell MS. 2007. Programming of body composition by early growth and nutrition. *Proc Nutr Soc* 66:423–434.
- Wells JC, Gigante DP, Wright A, Hallal PC, Victora CG. 2003. Validation of leg-to-leg impedance for body composition assessment in male Brazilians aged 16–19 years. *Int J Body Comp Res* 1:1–6.
- Wells JC, Hallal PC, Wright A, Singhal A, Victora CG. 2005. Fetal, infant and childhood growth: relationships with body composition in Brazilian boys aged 9 years. *Int J Obes (Lond)* 29:1192–1198.
- Winter JS, Hughes IA, Reyes FI, Faiman C. 1976. Pituitary-gonadal relations in infancy. II. Patterns of serum gonadotropin concentrations from birth to two years of age. *J Clin Endocrinol Metab* 42:679–686.
- Yajnik CS. 2002. The lifecycle effects of nutrition and body size on adult adiposity, diabetes and cardiovascular disease. *Obes Rev* 3:217–224.
- Yajnik CS. 2004. Early life origins of insulin resistance and type 2 diabetes in India and other Asian countries. *J Nutr* 134:205–210.